



National Library  
of Medicine

PubMed

PubMed

Nucleotide

Protein

Genome

Structure

PopSet

Search

PubMed



for

Go

Clear

Limits

Index

History

Clipboard

Display

Abstract



Save

Text

Order

Add to Clipboard

Entrez PubMed

☐ 1 : *Methods Enzymol* 1989;169:104-17

[Related Articles, Books](#)

PubMed Services

**Platelet interaction with artificial surfaces: in vitro evaluation.**

Lindon JN, Kushner L, Salzman EW

PMID: 2716522, UI: 89237928

Related Resources

Display

Abstract



Save

Text

Order

Add to Clipboard

Revised: January 10, 2000.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

National Library  
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	
Search PubMed	for				Go	Clear
<a href="#">Limits</a> <a href="#">Index</a> <a href="#">History</a> <a href="#">Clipboard</a>						

Entrez PubMed

☐ 1 : *J Am Acad Dermatol* 1990 May;22(5 Pt 1):781-91 [Related Articles, Books](#)

PubMed Services

**The effect of platelet releasate on wound healing in animal models.****Ksander GA, Sawamura SJ, Ogawa Y, Sundsmo J, McPherson JM**

Celtrix Laboratories, Collagen Corporation, Palo Alto, CA 94303.

Related Resources

The alpha granules of platelets contain growth factors that are important in wound healing. We found that a major effect of thrombin-induced human platelet releasates in animal models of wound healing is to enhance the development of granulation tissue and new connective tissue matrix. These studies provide further evidence that platelet-derived protein factors may be useful in treating full-thickness dermal wounds by increasing the rate of granulation tissue formation.

PMID: 2347964, UI: 90270507

Display	Abstract	Save	Text	Order	Add to Clipboard
---------	----------	------	------	-------	------------------

Revised: January 10, 2000.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)



National Library  
of Medicine

PubMed

PubMed Nucleotide Protein Genome Structure PopSet  
Search PubMed ☒ for  Go Clear  
☐ Limits Index History Clipboard

Entrez PubMed

☐ 1 : *Ann N Y Acad Sci* 1990;593:135-47

[Related Articles, Books](#)

PubMed Services

**Transforming growth factors-beta 1 and beta 2 enhance connective tissue formation in animal models of dermal wound healing by secondary intent.**

**Ksander GA, Chu GH, McMullin H, Ogawa Y, Pratt BM, Rosenblatt JS, McPherson JM**

Related Resources

Celtrix Laboratories, Collagen Corporation, Palo Alto, California 94303.

PMID: 2375592, UI: 90328644

Display Abstract ☐ Save Text Order Add to Clipboard

Revised: January 10, 2000.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)



National Library  
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet
Search PubMed	<input checked="" type="checkbox"/> for				Go Clear
<input type="checkbox"/> Limits Index History Clipboard					

Entrez PubMed

☐ 1 : *Am J Surg* 1993 Jul;166(1):74-81

[Related Articles, Books](#)

PubMed Services

## Growth factors and wound healing: Part II. Role in normal and chronic wound healing.

Bennett NT, Schultz GS

Department of Surgery, University of Florida, Gainesville.

Related Resources

Wound healing is a complex biologic process that involves the integration of inflammation, mitosis, angiogenesis, synthesis, and remodeling of the extracellular matrix. Part II of this two-part series reviews the results of experiments that indicate that growth factors and their receptors regulate key aspects of soft and hard tissue repair. Results of clinical studies are also reviewed that demonstrate that growth factor treatment accelerates healing of normal tissues and promotes healing of impaired wounds.

Publication Types:

- Review
- Review, tutorial

PMID: 8392302, UI: 93319056

Display	Abstract	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------------	------	------	-------	------------------

Revised: January 10, 2000.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)

National Library  
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	
Search PubMed	for				Go	Clear
<a href="#">Limits</a> <a href="#">Index</a> <a href="#">History</a> <a href="#">Clipboard</a>						
Display Abstract <input checked="" type="checkbox"/> Save Text Order Add to Clipboard						

Entrez PubMed

☐ 1 : *Am J Surg* 1993 Jun;165(6):728-37[Related Articles, Books](#)

PubMed Services

**Growth factors and wound healing: biochemical properties of growth factors and their receptors.****Bennett NT, Schultz GS**

Department of Surgery, University of Florida, Gainesville.

Related Resources

Wound healing is a complex biologic process that involves chemotaxis and division of cells, neovascularization, synthesis of extracellular matrix proteins, and remodeling of scar. Peptide growth factors have been shown to regulate many of these processes in vitro, leading to the hypothesis that peptide growth factors also regulate important phases of wound healing in vivo. Part I of this two-part series presents an overview of the biochemical properties of five families of peptide growth factors that are thought to be involved in wound healing: epidermal growth factor (EGF), transforming growth-factor-beta (TGF-beta), platelet-derived growth factor (PDGF), insulin-like growth factor (IGF), and fibroblast growth factor (FGF).

**Publication Types:**

- Review
- Review, academic

PMID: 8506974, UI: 93282541

Display	Abstract	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------------	------	------	-------	------------------

Revised: January 10, 2000.

[Disclaimer](#) | [Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)